Maryland COVID-19 Vaccination Plan



Vaccine	Туре	How It Works	Dosing	Possible Side Effects
Pfizer / BioNTech	mRNA	Contains genetic material that instructs the immune system to build immunity via Tlymphocytes, B-lymphocytes, and neutralizing antibodies	Two doses, delivered three weeks apart.	Injection-site pain, fatigue, headaches, chills.
Moderna			Two doses, delivered four weeks apart.	Injection-site pain, fatigue, muscle aches, joint pain, headaches, chills.
Johnson & Johnson	Vector	Contains a genetically modified version of a non-replicating adenovirus as a viral vector for coronavirus spike protein DNA which then prompts the body's immune system to activate T-lymphocytes and antibodyproducing B-lymphocytes to attack infected cells and induce immunity.	One dose	Injection site pain, fatigue, headache, muscle aches, joint pain
AstraZeneca			Two doses, delivered four weeks apart. NOTE: A half-dose may be recommended for the first dose.	Fever, muscle aches, headaches, injection- site pain.
Novavax	Protein Subunit	Uses synthetic coronavirus spike proteins to activate the body's immune system to produce antibodies against COVID-19, as well as an adjuvant to boost its effects.	Two doses, delivered three weeks apart.	Injection site pain, rash, headaches, muscle aches, fever

What You Need To Know About The Vaccines

- People who have recovered from COVID-19 may still benefit from getting vaccinated.
- COVID-19 vaccines will not alter the body's chemistry or DNA, and they will not cause patients to test positive.
- Because of limited supply initially, some groups will be recommended to get a COVID-19 vaccine before others.
 Supply is expected to increase greatly in the future.
- Vaccinated patients may still transmit COVID-19—but as greater numbers of people are vaccinated, we will achieve "herd immunity" that will greatly reduce cases of serious illness and death from COVID-19.

Key Facts For Your Patients

The vaccines are safe, and they will not give you COVID-19.

The vaccines are effective at preventing you from getting seriously ill with COVID-19.

Mild side effects are normal, including injection-site pain, fatigue, fever, aches, and chills.